



West African Journal of Life Sciences. Available online at <http://www.wajls.com> ISSN (Print): 2992-5150. Volume 2; Issue 1: pages 10-23

Full Length Research

Effects of Preservation Methods on Nucleic Acid Yield and Integrity from African Yam Bean (*Sphenostylis Stenocarpahochst. Ex A. Rich.*) Tissues for Sustainable Molecular Research

Lawani, M^{1*} and Shittu, H²

¹Department of Plant Science and Biotechnology - Faculty of Life Sciences, Ambrose Alli University, Ekpoma, Nigeria.

²Department of Plant Biology and Biotechnology - Faculty of Life Sciences, University of Benin, Edo State, Nigeria.

*Corresponding Author's Email: lawani.mathew@aauekpoma.edu.ng, Tel +234 810 453 9058

ABSTRACT

The African yam bean (AYB) (*Sphenostylis stenocarpa* Hochst. Ex A. Rich.) is an underutilized legume native to sub-Saharan Africa, valued for its edible seeds and tubers rich in proteins, carbohydrates, and essential micronutrients. This study aimed to optimize cost-effective preservation methods for African Yam Bean leaf tissues to ensure high-quality Deoxyribonucleic acid extraction, critical for molecular biology applications. African Yam Bean leaves were subjected to various preservation conditions, including chemical buffers (1× CTAB, 1× TAE, and 70% ethanol) and temperature settings (-20°C, 4°C, 25°C, and >25°C). Results indicated that freezing at -20°C yielded the highest Deoxyribonucleic acid (DNA) purity and stability, with consistent A₂₆₀/A₂₈₀ ratios (1.7–2.0). Ethanol preservation also demonstrated strong performance, providing a viable alternative for resource-constrained environments. The DNA preserved with cetyltrimethylammonium bromide and Tris-acetate-ethylenediaminetetraacetic acid buffers exhibited lower stability and increased contamination over time. These findings highlight practical and scalable methods to enhance DNA integrity from plant tissues, advancing the utility of AYB in molecular biology research and sustainable agriculture.

Received
February 24,
2025

Received in
Revised form,
May 31, 2025

Accepted June
3, 2025

Available
Online: June
12, 2025

W. Afr. J. Life
Sci. 2(1):10-23

Keywords: African yam bean, DNA extraction, molecular biology, preservation methods, sustainable agriculture.

INTRODUCTION

The African yam bean (*Sphenostylis stenocarpa*) is a nutritionally rich but underutilized legume native to sub-Saharan Africa. The African yam bean belongs to the family Leguminosae (Fabaceae), an annual, climbing or prostrate vine (Plate 1). This resilient crop produces both edible seeds and tubers and can be found in different varieties as shown in Plate 2. The crop is a valuable source of protein, carbohydrates, and micronutrients such as iron and zinc. Despite its high nutritional potential, the crop remains marginalized due to limited awareness, inadequate breeding programs, and its association with traditional, subsistence-level farming (Gbenga-Fabusiwa, 2021; Baiyeri et al., 2018).

The African yam bean is highly adaptable, thriving in diverse and often marginal soils with minimal inputs, making it a promising candidate for improving food security in resource-constrained regions (Oagile et al., 2012). Its seeds are comparable to other legumes in nutritional content, while the tubers provide additional dietary energy, creating a dual-purpose crop with significant potential for food diversification (Adewale & Nnamani, 2022). This crop is also recognized for its contributions to sustainable agriculture. African yam bean nodulates with rhizobia, enhancing nitrogen fixation and reducing dependency on synthetic fertilizers (Assefa & Kleiner, 1997).

However, its broader adoption is hindered by challenges such as limited processing knowledge, anti-nutritional factors, and insufficient research into its genetic diversity and agronomic practices (Gbenga-Fabusiwa, 2021). In light of global efforts to combat malnutrition and diversify food systems, the African yam bean offers a unique

opportunity to integrate an overlooked crop into modern agricultural and nutritional strategies. Its potential for genetic improvement, coupled with its adaptability and nutritional value, makes it a promising candidate for addressing the dual challenges of food security and sustainable farming in the face of climate change (Shitta et al., 2021).



Plate. 1. African Yam Bean




Plate. 2. Varieties of African Yam Bean seeds (Gbenga-Fabusiwa, 2021)

An oval shaped cream colour variety with black eye colour (A) A reddish brown variety with black eye colour (B)

Nucleic acid (NA) extraction is a cornerstone of molecular biology, enabling a wide array of applications in research, diagnostics, and therapeutic development. The quality and purity of extracted Deoxyribonucleic acid (DNA) and Ribonucleic acid (RNA) are critical for the accuracy of downstream molecular techniques, including polymerase chain reaction (PCR), next-generation sequencing (NGS), and gene expression analysis (Widen & Silbert, 2016). Emerging technologies, such as automated and portable systems, are making nucleic acid extraction more accessible. These innovations are critical for point-of-care diagnostics, enabling rapid, efficient preparation of nucleic acids even in resource-limited settings (Paul et al., 2020). Extraction of high-quality DNA in terms of quality and quantity is necessary for molecular biology studies. Generally, samples are obtained fresh and their DNA extracted for research purposes, but in a situation where the place of sample collection is far from the laboratory, the need for preservation therefore arises.

Preservation methods are complicated and expensive and its apparatus is hard to come by, as well as the possibility of health hazards issues developing from them. For instance - 80°C freezer is very expensive and not readily found in institutions in developing countries like Nigeria. Naphthalene, a preservative is both carcinogenic and dangerous to the eyes; liquid nitrogen which can be used to freeze-dry plant materials is volatile and needs optimal direction when used. There is therefore a need for alternative preservation methods. Works of literature have shown that silica gel desiccation of leaf

samples and immersion in NaCl-Cetyltrimethylammonium bromide (CTAB) solutions have been effective. Quality DNA extracts were obtained from tissues preserved for over a month using these methods (Chase & Hills, 1991; Bhattacharjee et al., 2009). Recently Johnson et al., 2023 reported that leaf samples preserved in ethanol, particularly 96% ethanol, demonstrate superior DNA quality but the procedure was combined with proteinase digestion.

This research aimed to develop cost-effective preservation methods for plant samples before DNA extraction, focusing on alternatives to those commonly reported in the literature. By addressing the challenges of implementing sophisticated and expensive techniques in developing countries, this study seeks to offer a broader range of practical options for researchers in molecular biology.

MATERIALS AND METHODS

Source of Plant Materials

African yam bean (*Sphenostylis stenocarpa* Hochst. Ex A. Rich.) leaves were collected from the Botanical Garden of the Department of Plant Biology and Biotechnology, Faculty of Life Sciences, University of Benin, Benin City, Edo State, Nigeria. They were authenticated at the departmental herbarium by the curator and assigned an ID number of UNIBEN/PSBBG/010624.

Chemicals and Reagents

The preservation buffers used were obtained from the Nigerian Institute for Oil Palm Research (NIFOR), Nigeria. They include 1× CTAB, 1× Tris-acetate-ethylenediaminetetraacetic acid, and 70% ethanol. The DNA extraction buffer was prepared as 10× CTAB, comprising 0.5 M Tris, 5 M NaCl, 0.3 M

ethylenediaminetetraacetic acid, 10% CTAB, and 20 g Polyvinylpyrrolidone (PVP). Additional reagents included chloroform: phenol (1:1), ice-cold 100% ethanol and 70% ethanol as precipitation buffers, sterile water for dissolving DNA, and TAE buffer (0.89 M Tris, 0.89 M acetic acid, 25 mM EDTA, pH 8.3) as a loading buffer. Ethidium bromide was used as the staining dye, and agarose was utilized for gel electrophoresis.

Sample Preservation

After collection, the leaves were preserved under various conditions to study their effects on DNA integrity. Specifically, preservation was carried out at four temperature settings (-20°C, 4°C, 25°C, and above 25°C in sunlight) and in three chemical buffers (1× CTAB, 1× TAE, and 70% ethanol). The samples were stored for durations of 3, 6, 9, and 12 days. Leaves from each condition were retrieved at these intervals and prepared for DNA extraction.

DNA Extraction

DNA extraction was performed using a CTAB-based protocol adapted from NACGRAB. Preserved leaf tissue of 0.3g was ground with 2 mL of preheated 10× CTAB buffer and 20 mg of acid-washed sand using a mortar and pestle, which were preheated at 65°C. The resulting homogenate was transferred into microfuge tubes and incubated at 65°C for 15 minutes in a water bath. The suspension was centrifuged at 12,500 rpm for 5 minutes at 4°C, and the supernatant was carefully transferred into a fresh tube. Chloroform: phenol (1:1) was added to the supernatant, mixed thoroughly, and centrifuged at 14,000 rpm for 5 minutes at 4°C. The aqueous phase was transferred to a new tube, and DNA was precipitated by adding 0.8 mL of ice-cold 100% ethanol and incubating the mixture at -20°C for 1 hour.

After centrifugation at 13,000 rpm for 20 minutes at 4°C, the pellet was washed three times with 500 µL of 70% ethanol, followed by centrifugation at 1,000 rpm for 3 minutes at 4°C. The DNA pellet was air-dried and dissolved in 100 µL of sterile water.

Spectrophotometric Analysis

DNA purity and yield were determined using Ultraviolet Spectrophotometry. To blank the spectrophotometer, 1.5 mL of sterile water was used in a cuvette. For analysis, 20 µL of each DNA extract was diluted in 1.48 mL of sterile water, and absorbance readings were taken at 260 nm and 280 nm. Triplicate readings were recorded for each sample. DNA purity was assessed using the A_{260}/A_{280} ratio, with values between 1.7 and 2.0 considered pure. The applied purity keys were: 1.7–2.0 for pure DNA, <1.7 for protein contamination, and >2.0 for RNA contamination.

DNA concentration was calculated using the formula

Concentration (ng/µL) = $A_{260} \times \text{Dilution Factor} \times 50$

Where A_{260} represents absorbance at 260 nm, the Dilution Factor accounts for sample dilution, and 50 is a standard conversion factor for double-stranded DNA. The DNA yield was determined by multiplying the concentration by the total volume of the extract and dividing by 1000 to convert from nanograms (ng) to micrograms (µg).

Agarose Gel Electrophoresis

To verify the integrity of the extracted DNA, agarose gel electrophoresis was performed. A 0.8% agarose gel was prepared by dissolving 0.8g of agarose in 100mL TAE buffer, followed by the addition of 1 µL of ethidium bromide. The gel was carefully poured into a tray with a well comb and allowed to set and

be cooled at room temperature for 20 minutes. DNA samples (8 μ L) were mixed with 2 μ L of loading dye, loaded into the gel wells, and subjected to electrophoresis under an electric current at 80V for 1 hour. DNA bands were visualized under UV light, and their clarity and consistency were assessed to determine DNA integrity.

STATISTICAL ANALYSIS

Significant differences between groups were assessed using Analysis of Variance (ANOVA) with a significance threshold of $p < 0.05$. The yields were averages calculated from triplicate experiments, with error margins reflecting the standard error of the mean (SEM).

RESULTS

Nucleic Acid Yield from Different Preservation Methods

Chemical Preservation

The nucleic acid yield was monitored across twelve (12) days using three chemical preservation methods, which were the preservation of African yam bean leaf tissues in 70% Ethanol, 1 $\times$ TAE, and 1 $\times$ CTAB. Nucleic acid yield under chemical preservation varied significantly across the methods with a threshold of $p < 0.05$.

Table 1 shows the nucleic acid yield in μ g/g from the chemical preservation method while Figure 1 presents a graphical representation of DNA yields obtained from the same preservation process. The 70% Ethanol preservation method showed consistent yields initially, but a marked decline on Day 12. The 1 $\times$ TAE preservation method maintained stable yields until Day 9, followed by a decline, while 1 $\times$ CTAB

showed fluctuating yields, peaking significantly on Day 9.

Table 1: Nucleic Acid Yield from Chemical Preservation Method

Time (Day)	70% Ethanol (μ g/g)	1 $\times$ TAE (μ g/g)	1 $\times$ CTAB (μ g/g)
3	20 $\pm$ 1.5	44 $\pm$ 3.2	13 $\pm$ 0.9
6	33 $\pm$ 2.1	47 $\pm$ 2.8	8 $\pm$ 0.6
9	59 $\pm$ 3.8	28 $\pm$ 2.0	126 $\pm$ 6.4
12	-2 $\pm$ 0.2	-3 $\pm$ 0.3	12 $\pm$ 0.8





Figure 1: Trends in Nucleic Acid Yield (Chemical Preservation).

Temperature Preservation

Temperature preservation revealed that -20 $^{\circ}$ C consistently maintained the highest yield, though there was variability on Day 6. Preservation at higher temperatures (>25 $^{\circ}$ C) and 4 $^{\circ}$ C showed reduced yields and lower stability over time. Table 2 shows the results obtained for nucleic acid yield isolated from AYB leaf tissues preserved in different temperature conditions for 3, 6, 9 and 12 days while Figure 2 is a graphical visualization of the various nucleic acid yields obtained from tissues preserved in the same temperature preservation methods and days.

Table 2: Nucleic Acid Yield from Temperature Preservation Method

Time (Day)	-20°C (µg/g)	4°C (µg/g)	25°C (µg/g)	>25°C (µg/g)
3	87 ± 5.4	4 ± 0.3	58 ± 3.8	31 ± 2.1
6	5 ± 0.5	22 ± 1.7	38 ± 2.9	58 ± 3.2
9	70 ± 4.2	55 ± 3.3	170 ± 8.5	125 ± 6.2
12	45 ± 3.2	40 ± 2.8	20 ± 1.4	16 ± 0.9

**Figure 2: Trends in Nucleic Acid Yield (Temperature Preservation)**

Nucleic Acid Purity across Preservation

Methods

The purity of DNA was evaluated using the A_{260}/A_{280} ratio, which indicated contamination by proteins or RNA.

>2.0) over the observation period. Table 3 shows the effect of three chemical treatments on nucleic acid purity for 3, 6, 9, and 12 days.

Chemical Preservation

DNA purity varied across different preservation methods, with a noticeable decline by Day 12. The use of 1× TAE demonstrated inconsistent purity levels, with evidence of both protein contamination (ratio <1.7) and RNA contamination (ratio

Table 3: Nucleic Acid Purity Ratios for Chemical Preservation

Time (Day)	70% Ethanol	1× TAE	1× CTAB
3	2.0 (Pure)	0.2 (<1.7, Protein)	1.7 (Pure)
6	2.4 (>2.0, RNA)	0.6 (<1.7, Protein)	2.1 (>2.0, RNA)
9	1.8 (Pure)	1.8 (Pure)	1.4 (<1.7, Protein)
12	- (No Data)	- (No Data)	2.8 (>2.0, RNA)


**Figure 3: Purity Trends in Chemical Preservation Method**

Figure 3 shows that pure DNA accounts for the largest proportion of samples, followed by RNA and protein contamination in nearly equal amounts. RNA contamination represents a significant fraction, indicating frequent co-extraction of RNA alongside DNA.

Temperature Preservation

DNA stored at -20°C consistently retained high purity across all time points, with A_{260}/A_{280} ratios remaining within the optimal range of 1.7–2.0. Conversely, elevated temperatures, especially $>25^{\circ}\text{C}$, led to increased RNA contamination, as indicated by ratios exceeding 2.0. Table 4 shows the effect of the temperature preservation method on nucleic acid purity for varying numbers of days.

Table 4: DNA Purity Ratios for Temperature Preservation

Time (Day)	-20°C	4°C	25°C	>25°C
3	1.8 (Pure)	1.6 (<1.7, Protein)	1.9 (Pure)	1.4 (<1.7, Protein)
6	1.9 (Pure)	2.3 (>2.0, RNA)	1.8 (Pure)	2.2 (>2.0, RNA)
9	2.0 (Pure)	1.9 (Pure)	1.5(<1.7, Protein)	1.8 (Pure)
12	2.0 (Pure)	1.9 (Pure)	2.8 (>2.0, RNA)	3.0 (>2.0, RNA)


**Figure 4: Purity Trends in Temperature Preservation Method.**

Figure 4 shows that pure DNA is the most prevalent, indicating that optimal storage in temperature conditions can effectively maintain nucleic acid integrity.

DNA Integrity Check by Gel Electrophoresis

This method was selected due to the specificity of ethidium bromide staining for DNA, enabling clear visualization of nucleic

acid bands. Figure 5 below illustrates the DNA profiles obtained from samples preserved under different conditions at intervals of 3, 6, 9, and 12 days. Each lane represents one of the four temperature conditions (arranged in the order: -20°C, 4°C, 25°C, and >25°C) and one of the three chemical preservation methods (arranged in the order: 1× CTAB, 1× TAE, and 70% ethanol).




Figure 5: Agarose electrophoresis gel images of nucleic acid extracts from preserved tissues of African yam bean (*Sphenostylis stenocarpa*).

Samples preserved for 3 days (A) 6 days (B) 9 days (C) and 12 days (D) providing a comparative view of nucleic acid integrity over time under different preservation conditions.

DISCUSSION

This study provides an in-depth evaluation of preservation methods for African yam bean (AYB) leaf tissues, contributing to a growing body of knowledge on sample preparation for molecular biology. The results confirmed that both the preservation method and duration had a significant impact on nucleic acid yield and purity. The findings indicate that chemical preservation methods demonstrated varying effectiveness in maintaining DNA yield and purity over time. Ethanol preservation proved highly effective initially, maintaining DNA integrity with minimal contamination.

However, DNA yield declined significantly by Day 12. This result aligns with Johnson et al. (2023) and Bressan et al. (2014) who highlighted the effectiveness of ethanol for preserving recalcitrant plant species, though their methodology incorporated additional proteinase digestion, suggesting a potential refinement for ethanol-based protocols. The 1× TAE buffer demonstrated stable yields

until Day 9, after which a decline was observed. Tris-based buffers like TAE have been reported to interact with DNA, altering its migration properties and potentially leading to instability (Stellwagen et al., 2000). The 1× CTAB method exhibited fluctuations, peaking significantly on Day 9. This transient peak observed in CTAB-preserved samples on Day 9 might reflect initial stabilization by its detergent properties, which degraded over time, leading to renewed enzymatic activity and reduced DNA stability. This aligns with findings that CTAB enhances DNA integrity initially by binding to nucleic acids and reducing nuclease activity but can later cause DNA shearing or contamination due to residual chemicals (Guertler et al., 2013). The decline in DNA yield with CTAB and TAE buffers could be linked to their inability to adequately inhibit enzymatic activity over extended durations. Additionally, these variations suggest that buffer selection is critical for maintaining nucleic acid integrity (Carey et al., 2023).

Temperature preservation results showed that -20°C consistently maintained the highest yield and DNA stability over time, with some variability on Day 6 which may be due to differences in sample handling, pipetting errors, or variations in extraction efficiency on that particular day. In addition, samples frozen over time may experience phase separation or ice layering, leading to uneven distribution of nucleic acids within the sample. This could make DNA extraction inconsistent on different days (Cordsmeier & Hahn, 2022). The superior performance of -20°C freezing in preserving DNA integrity likely comes from its ability to arrest enzymatic activities and stabilize macromolecular structures, thereby minimizing degradation. Studies have shown that freezing at -20°C maintains DNA yield and quality over extended periods, comparable to fresh samples, by preventing enzymatic hydrolysis and oxidative damage (Wood & Wang, 2024).

However, repeated freeze-thaw cycles have been reported to cause progressive DNA degradation, particularly for high-molecular-weight fragments (Shao et al., 2012). Furthermore, Bainard et al. (2010) reported that freezing at -20°C effectively preserved DNA integrity across diverse plant tissues. Preservation at higher temperatures ($>25^{\circ}\text{C}$) and 4°C resulted in reduced yields and increasing contamination. These observations align with Paul et al. (2020), who demonstrated that high temperatures accelerate nucleic acid degradation through enzymatic and oxidative pathways.

The work of Michaud and Foran (2011) also supports these findings for preservation in lower and higher temperatures, showing that refrigerated storage was intermediate in

effectiveness while desiccation was least effective for long-term preservation. DNA purity evaluation showed that the chemical preservation methods exhibited varying DNA purity levels. Ethanol generally maintained purity but showed a noticeable decline by Day 12. Ethanol's efficacy in DNA preservation may be attributed to its ability to dehydrate the cellular matrix, effectively halting enzymatic functions responsible for DNA breakdown.

Ethanol has been demonstrated to maintain DNA stability by preventing hydrolytic degradation and protein-DNA interactions, particularly in plant and insect specimens (Marquina et al., 2021). The $1\times$ TAE method exhibited inconsistent purity levels, with evidence of protein contamination ($A_{260}/A_{280} < 1.7$) and RNA contamination ($A_{260}/A_{280} > 2.0$) over time. The $1\times$ CTAB method also showed fluctuations, with protein and RNA contaminations becoming evident at later time points. The results suggest that the peak in terms of DNA yield observed on Day 9 from samples preserved in $1\times$ CTAB were contaminated by protein and therefore not fit for downstream experiments (Guertler et al., 2013).

DNA stored at -20°C consistently retained high purity, with A_{260}/A_{280} ratios remaining within the optimal range of 1.7–2.0 across all time points. In contrast, elevated temperatures ($>25^{\circ}\text{C}$) resulted in increased RNA contamination, as indicated by ratios exceeding 2.0. This is in line with Sadler & Khodavirdi (2015), who demonstrated that viable RNA could be extracted from tissue samples stored at room temperature for up to three months, challenging the assumption that RNA rapidly degrades at high temperatures. The results further suggest that higher temperatures may promote RNA

retention rather than degradation under certain conditions. Protein contamination, though slightly less frequent, remained a concern, implying that some preservation methods may not sufficiently inhibit protein carryover (McNevin, 2016).

Overall, Figures 3 and 4 showed that the temperature preservation method yielded purer DNA (41.1%) compared to chemical preservation (40%). The distribution highlights the influence of temperature on nucleic acid purity and the need for precise storage conditions to minimize contamination. This study also highlights that preservation methods influence not only the quantity of extracted DNA but also its usability for downstream applications. Freezing and ethanol preservation produced DNA with high purity, suitable for sensitive molecular techniques like PCR and sequencing, whereas DNA preserved with CTAB and TAE exhibited contamination, potentially limiting its use.

CONCLUSION

This study identifies freezing at -20°C as the most effective preservation method for African yam bean leaf tissues in terms of DNA yield and purity, consistently producing DNA suitable for molecular applications. Ethanol preservation also demonstrated excellent performance, providing a viable, cost-effective alternative for resource-limited environments. In contrast, preservation using CTAB and TAE buffers showed limitations, with reduced DNA stability over time. These findings provide a practical framework for enhancing

While the study establishes the efficacy of specific preservation methods, some limitations require consideration. The influence of secondary metabolites in AYB leaves on DNA extraction efficiency remains an open question. These metabolites might interact with preservation agents, influencing yield and purity. Additionally, the variability introduced by manual handling during preservation and extraction, though minimized through standard protocols, could be further refined with automated processes.

Future research should explore testing these methods across other underutilized and recalcitrant species to validate their efficacy. Another area should be in investigating the interactions between plant metabolites and preservation chemicals to improve DNA yield and purity.

molecular biology preservative research in developing countries and contribute to the broader goal of integrating underutilized crops into sustainable agricultural systems.

ACKNOWLEDGEMENTS

I sincerely acknowledge my supervisor, Prof. Shittu Hakeem, for his invaluable guidance, leadership, and generous investment of time throughout this research. His insightful feedback, unwavering support, and academic mentorship were instrumental to the success of this work.

REFERENCES

- Adewale, B. D., & Nnamani, C. V. (2022). Introduction to food, feed, and health wealth in African yam bean, a locked-in African indigenous tuberous legume. *Frontiers in Sustainable Food Systems*, 6,

726458.

<https://doi.org/10.3389/fsufs.2022.726458>

- Assefa, F., & Kleiner, D. (1997). Nodulation of African yam bean (*Sphenostylis stenocarpa*) by *Bradyrhizobium* sp.

- isolated from *Erythrina brucei*. *Biology and Fertility of Soils*, 25, 209–210. <https://doi.org/10.1007/s003740050305>
- Bainard, L. D., Klironomos, J. N., & Hart, M. M. (2010). Differential effect of sample preservation methods on plant and arbuscular mycorrhizal fungal DNA. *Journal of Microbiological Methods*, 82(2), 124–130. <https://doi.org/10.1016/j.mimet.2010.05.001>
- Baiyeri, S. O., Uguru, M. I., Ogbonna, P. E., Samuel-Baiyeri, C. C. A., Okechukwu, R., Kumaga, F. K., & Amoatey, C. (2018). Evaluation of the nutritional composition of the seeds of some selected African yam bean (*Sphenostylis stenocarpa* Hochst Ex. A. Rich (Harms)) accessions. *Agro-Science*, 17(2), 37–44. <https://doi.org/10.4314/as.v17i2.5>
- Bhattacharjee, R., Ferguson, M., Gedil, M., Dumet, D., & Ingelbrecht, I. (2009). Field collection, preservation and large scale DNA extraction procedures for cassava (*Manihot esculenta* Crantz.). *African Journal of Biotechnology*, 8(15), 3424–3430. <https://doi.org/10.5897/AJB2009.000-9334>
- Bressan, E. A., Rossi, M. L., Gerald, L. T., & Figueira, A. (2014). Extraction of high-quality DNA from ethanol-preserved tropical plant tissues. *BMC Research Notes*, 7, 268. <https://doi.org/10.1186/1756-0500-7-268>
- Carey, S. J., Becklund, L. E., Fabre, P. P., & Schenk, J. J. (2023). Optimizing the lysis step in CTAB DNA extractions of silica-dried and herbarium leaf tissues. *Applications in Plant Sciences*, 11(3), e11522. <https://doi.org/10.1002/aps3.11522>
- Chase, M. W., & Hills, H. H. (1991). Silica gel: An ideal material for field preservation of leaf samples for DNA studies. *Taxon*, 40(2), 215–220. <https://doi.org/10.2307/1222975>
- Cordsmeier, L., & Hahn, M. B. (2022). DNA stability in biodosimetry, pharmacy and DNA based data-storage: Optimal storage and handling conditions. *ChemBioChem*, 23(20), e202200391. <https://doi.org/10.1002/cbic.202200391>
- Gbenga-Fabusiwa, F. J. (2021). African yam beans (*Sphenostylis stenocarpa*): A review of a novel tropical food plant for human nutrition, health and food security. *African Journal of Food Science*, 15(2), 33–47. <https://doi.org/10.5897/AJFS2020.1961>
- Guertler, P., Harwardt, A., Eichelinger, A., Muschler, P., Goerlich, O., & Busch, U. (2013). Development of a CTAB buffer-based automated gDNA extraction method for the surveillance of GMO in seed. *European Food Research and Technology*, 236, 599–606. <https://doi.org/10.1007/s00217-013-1916-y>
- Haile, A. T., Johnsen, S. S., Kovi, M. R., Hvoslef-Eide, T., Tesfaye, B., & Rognli, O. (2021). Comparison of different leaf preservation methods to obtain high-quality DNA from enset. *PeerJ*, 9, e11719. <https://doi.org/10.7717/peerj.11719>
- Johnson, G., Canty, S. W. J., Lichter-Marck, I. H., Wagner, W. L., & Wen, J. (2023). Ethanol preservation and pretreatments facilitate quality DNA extractions in recalcitrant plant species. *Applications in Plant Sciences*, 11(3), e11519. <https://doi.org/10.1002/aps3.11519>
- Marquina, D., Buczek, M., Ronquist, F., & Łukasik, P. (2021). The effect of ethanol concentration on the morphological and molecular preservation of insects for biodiversity studies. *PeerJ*, 9, e10799. <https://doi.org/10.7717/peerj.10799>
- McNevin, D. (2016). Preservation of and DNA extraction from muscle tissue. In J. R. Lakshmanan (Ed.), *Methods in Molecular Biology* (Vol. 1420, pp. 43–53). Humana Press. https://doi.org/10.1007/978-1-4939-3600-7_4
- Michaud, C. L., & Foran, D. R. (2011). Simplified field preservation of tissues for subsequent DNA analyses. *Journal of Forensic Sciences*, 56(4), 846–852. <https://doi.org/10.1111/j.1556-4029.2011.01771.x>

- Oagile, O., Mmolotsi, R., Segwagwe, A., & Babili, T. P. (2012). African yam bean (*Sphenostylis stenocarpa*) nodulates promiscuously with *Rhizobium* indigenous to soils of Botswana. *African Journal of Biotechnology*, 11(24), 6510–6515.
<https://doi.org/10.5897/AJB11.1994>
- Paul, R., Ostermann, E., & Wei, Q. (2020). Advances in point-of-care nucleic acid extraction technologies for rapid diagnosis of human and plant diseases. *Biosensors and Bioelectronics*, 169, 112592.
<https://doi.org/10.1016/j.bios.2020.112592>
- Shao, W., Khin, S., & Kopp, W. C. (2012). Characterization of effect of repeated freeze and thaw cycles on stability of genomic DNA using pulsed field gel electrophoresis. *Biopreservation and Biobanking*, 10(1), 4–11.
<https://doi.org/10.1089/bio.2011.0016>
- Shitta, N. S., Abtew, W. G., Ndlovu, N., Oselebe, H. O., Edemodu, A. C., & Abebe, A. T. (2021). Morphological characterization and genotypic identity of African yam bean (*Sphenostylis stenocarpa* Hochst ex. A. Rich. Harms) germplasm from diverse ecological zones. *Plant Genetic Resources*, 19(1), 58–66.
<https://doi.org/10.1017/S1479262121000095>
- Shittu, H. O., Durojaye, A., Uzama, O., Odenore, V., & Eke, C. (2022). Optimizing DNA extraction without lyophilization or liquid nitrogen grinding from African yam bean (*Sphenostylis stenocarpa* Hochst Ex. A. Rich.) Harms tissues. *Nigerian Journal of Life Sciences*, 4(2), 192–198.
<http://dx.doi.org/10.52417/njls.v4i2.199>
- Stellwagen, N. C., Gelfi, C., & Righetti, P. G. (2000). DNA and buffers: The hidden danger of complex formation. *Biopolymers*, 54(2), 137–142.
[http://dx.doi.org/10.1002/1097-0282\(200008\)54:2%3C137::AID-BIP6%3E3.0.CO;2-I](http://dx.doi.org/10.1002/1097-0282(200008)54:2%3C137::AID-BIP6%3E3.0.CO;2-I)
- Straube, D., & Juen, A. (2013). Storage and shipping of tissue samples for DNA analyses: A case study on earthworms. *European Journal of Soil Biology*, 57, 13–18.
<https://doi.org/10.1016/j.ejsobi.2013.04.001>
- Wang, X., Li, L., Zhao, J., Li, F., Guo, W., & Chen, X. (2017). Effects of different preservation methods on inter simple sequence repeat (ISSR) and random amplified polymorphic DNA (RAPD) molecular markers in botanic samples. *Comptes Rendus Biologies*, 340(4), 204–213.
<https://doi.org/10.1016/j.crvi.2017.03.002>
- Widen, R. H., & Silbert, S. (2016). Nucleic acid extraction in diagnostic virology. In D. D. Richman, R. J. Whitley, & F. G. Hayden (Eds.), *Clinical Virology Manual* (5th ed., pp. 117–128). ASM Press.
<https://doi.org/10.1128/9781555819156.ch10>
- Wood, A., & Wang, Y. (2024). B-184 Purifying high-quality genomic DNA from frozen blood samples stored up to 1.5 years at –20°C. *Clinical Chemistry*, 70(Supplement_1), hvae106-544.
<https://doi.org/10.1093/clinchem/hvae106.544>